

Grapevine Berry Phenolic Compounds in Relation with Plant Water Status

Phenolische Substanzen in Weinbeeren in Abhängigkeit von der Wasserversorgung

Alain Deloire¹, Hernan Ojeda^{1,2,3} Brigitte Federspiel¹,
Elena Kraeva^{1,3}, Claude Andary³.

Grape derived secondary metabolites are the principal sources of wine aroma, flavor, color and taste. Most of available data about phenolics are concerned mature berries. Nevertheless, the contribution of flavonols in metabolism may be important, not only in red berries, but especially in green, when the berries are most susceptible to different kind of stresses. The changes in phenol composition in green berries, under water stress conditions (Ojeda *et al.*, 2001; in press) and with different training systems (Kraeva *et al.*, 2001; submitted) have been observed

We studied the black grapevine varieties *Vitis vinifera* L cv. Syrah and Grenache noir. The experiments were carried at different stages of grape growth in controlled conditions (adult plants grown in containers with a drip irrigation) and in the vineyard (terroir effect). The training systems were Lyre and Espalier pruned as cordon. The plant water status was estimated by predawn leaf water potential (table 1).

Table 1: Plant water status was estimated by predawn leaf water potential. Syrah are irrigated adult vines in pot : (T) = controlled vines; (S) = stressed vines. Grenache are vines in experimental vineyard : (NI) = non irrigated vines (terroir reference); (I) = irrigated vines in the same terroir.

Varieties	Predawn leaf water potential (Ψ_b) from anthesis to véraison	Predawn leaf water potential (Ψ_b) from véraison to maturity	
Syrah	$\leq -0,2$ Mpa	T	S
		$\leq -0,2$ Mpa	$\geq -0,6$ Mpa
Grenache	$\leq -0,3$ Mpa	I	NI
		$\pm -0,4$ Mpa	$\pm -0,6$ Mpa

¹ Agro.M., Viticulture-Oenologie, UMR «Sciences pour l'Oenologie», 2 place Viala, 34060 Montpellier. Corresponding author: e-mail: deloire@ensam.inra.fr

² EEA Mendoza INTA, San Martín 3853, 5507 Luján de Cuyo, Mendoza, Argentina.

³ Laboratoire de Botanique, Phytochimie et Mycologie, CNRS-UPR 9056, Faculté de Pharmacie, 15 av. C. Flahault, 34060 Montpellier, cédex 2, France.

The effects of plant water status on berry growth was studied (Ojeda *et al.*, 1998, 1999 and 2001). Water deficit did not affect cell division as we have shown by DNA extraction profiles. Reduction of berry size and berry weight was caused exclusively by a decrease of pericarp volume. Thus water deficit increased the skin-to-pulp ration, which has important consequence on both phenolic biosynthesis and concentration.

Water deficit level occurred during véraison to maturity period stimulate the flavonol biosynthesis (Figure 1). The training system, in relation with bunches microclimate, could modify the berry phenolic composition (Kraeva *et al.*, 2001; submitted).

The flavonol compounds are synthesised in the epidermal cells of berry skin and sequestered as various glycosides in the central vacuoles. Flavonols are synthesised in the epidermis. The quercetin (a major flavonol compound) is predominantly present in the glycosylated form in grape berries (Price *et al.*, 1995). It has been suggested that these glycosides act as UV screening compounds, helping to protect the plant tissue from damage, but also against pathogens (Dai *et al.*, 1995a Deloire *et al.*, 1998). Moreover, we have shown (Kraeva *et al.*, 2001, submitted) the role of flavans and flavonols against the elicitation with salicylic acid (SA), which was known to enhanced oxidative stress and induced defense reaction in plants (Chen *et al.*, 1993).




Figure 1: Flavonols content expressed in mg rutin equivalent, of Syrah berries of control plants and of plants affected by to water deficit treatment. C = control (100% ETP). S = water deficit between veraison and maturity (30% ETP). The arrow indicate the onset of veraison (softening of 10% of the berries).

Bibliographie

- Chen Z., Silva H., Klessig D.F. 1993 – Active oxygen species in the induction of plant systematic acquired resistance by salicylic acid. *Science*, 262, 1883.
- Dai G.H., Andary C, Mondolot-Cosson L, Boubals D. 1995a. Histochemical studies on the interaction between three species of grapevine, *Vitis vinifera*, *V. rupestris* and *V. rotundifolia* and the downy mildew fungus, *Plasmopara viticola*. *Physiol. Mol. Plant Pathol.*, 46, 177 - 188.
- Dai G. H., Andary C., Mondolot-Cosson L., Boubals D. 1995b - Histochemical responses of leaves of *in vitro* plantlets of *Vitis* spp. to infection with *Plasmopara viticola*. *Phytopathology*, 85, 149-154.
- Deloire A., Kraeva E., Dai G.H., Renault A.S., Rochard J., Chatelain C., Carbonneau A., Andary C. 1998. Les mécanismes de défense de la vigne. Des utilisations possibles pour lutter contre les pathogènes. *Phytoma*, 510, 46-51.
- Ojeda H., Andary C., Kraeva E., Carbonneau A., Deloire A. 2001 - Influence of pre and post-veraison water deficit on the synthesis and concentration of skin phenolic compounds during berry growth of *Vitis vinifera* L., cv Shiraz. *Am. J. of Enol. and Vitic* (in press).
- Ojeda H., Deloire A., Carbonneau A. 2001 – Influence of water deficits on grape berry growth. *Vitis* 40 (3), 141-145.
- Ojeda H., Deloire A., Carbonneau A., Ageorges A., Romieu A. 1999 – Berry development of grapevines : relations between the growth of berries and their DNA content indicate cell multiplication and enlargement. *Vitis* 38, 145-150.
- Ojeda H., Lebon E., Romieu C., Carbonneau A., Andary C., Deloire A. 1998 – Relations entre le stress hydrique et la croissance des baies de *Vitis vinifera* L. cv Syrah. Evolution des phénols. Gesco XI^{ème} journées du groupe européen d'étude des systèmes de conduite de la vigne, Sicile-Italie, 6-12 juin, 1, 185-192.
- Price S.H., Breen P.J., Valalladao M. and Watson .B.T. 1995 – Cluster sun exposure and quercetin in grapes and wine. *American Journal of Enology and Viticulture* 46, 187-194.