


# Symbiotic nitrogen fixation and N-transfer in a grass-white clover mixture





FINN P. VINTHER

Danish Institute of Agricultural Sciences, Dept. of Crop Physiology and Soil Science, Research Centre Foulum, DK-8830 Tjele, Denmark. E-mail: Finn.Vinther@agrsci.dk


This study constituted a part of a larger investigation, where input, losses, and recycling of nitrogen were estimated in either grass-clover or fertilised ryegrass grazed by dairy cows. The objective of the present study was to estimate the input of symbiotically fixed  $N_2$  into a grass-white clover mixture and the transfer of fixed  $N_2$  from clover to grass. The study included an evaluation of monoculture ryegrass and ryegrass in the mixture as reference crop for estimating SNF in grass-clover mixtures. Further, in a parallel experiment the amount of nitrogen incorporated after a 2<sup>nd</sup> year grass-clover were estimated.




Due to a cold and wet spring in 1994 the clover was poorly established and the total dry matter production corresponded to 7.5 tons ha<sup>-1</sup> compared to 10.6 tons ha<sup>-1</sup> in 1995. These differences also resulted in large difference in the amount of N fixed between the two years (see below).



There was no significant difference between the two methods determining the proportion of N derived from atmosphere (pNdfa). The proportion of grass N transferred from clover to grass (pgNdfa) tended to increase during the growing season.



About 50% of the nitrogen incorporated into the soil was estimated to be derived from  $N_2$  fixation.



The total  $N_2$  fixation corresponded to 160 kg N ha<sup>-1</sup> in 1994 and to 290 kg N ha<sup>-1</sup> in 1995.