

3rd ISOFAR Conference 2011

Use of competitive crops to reduce *Cirsium arvense*

Ilse A. Rasmussen

Formerly employed at Aarhus University

All photos courtesy of:

Henning C. Thomsen, Eugene Driessen, Bo Melander Aarhus University



From the weed garden of prof. Korsmo (http://foto.bioforsk.no/fotoweb/ugras/UK/Index.fwx)

Cirsium arvense *llse A. Rasmussen*

30th September, 2011

Cirsium arvense








Cirsium arvense Ilse A. Rasmussen

Biomass of C. arvense in different crops with and without catch crops/stubble cultivation




From Rasmussen

et al. 2005



C. arvense biomass on sandy loam in different crop types in two rotations





Crop rotation in experiment


Year	Field 1	Field 2	Field 3	Field 4
2004	Sugar beets	S. barley	Grass-clover	W. wheat + catch crop
2005	S. barley	Grass-clover	Potatoes	S. oats + catch crop
2006	Grass-clover	Potatoes	W. wheat + catch crop	S. barley
2007	Potatoes	W. wheat + catch crop	S. barley	Grass-clover
2008	W. wheat + catch crop	S. barley	Grass-clover	Potatoes
2009	W. oilseed rape	Grass-clover	Grass-clover	Hemp
2010	S. barley	S. barley	S. barley	S. barley








C. arvense number m⁻²





C. arvense biomass g m⁻²



Can competitive crops be used to reduce Cirsium arvense?

- > Competitive crops should be one of the strategies used in the management
- > A crop that is expected to be competitive, may sometimes not be so
- > The crops shown here are examples, others could be relevant in other situations.

Acknowledgements

This work was co-funded by

> The Foundation for Organic Farming (Fonden for Økologisk Landbrug) and the

> Danish Ministry of Food, Agriculture and Fisheries.

The author wishes to thank the technicians
Eugene Driessen and Karen Heinager
for their skilled and hard work in carrying out the experiment.