

HAGEL, I., D. BAUER, S. HANEKLAUS and E. SCHNUG (2000): *Quality Assessment of Summer and Autumn Carrots from a Biodynamic Breeding Project and Correlations of Physico-Chemical Parameters and Features Determined by Picture Forming Methods. Proceedings of the 13th IFOAM Scientific Conference, Basel, Switzerland, 284-287.*

## Quality Assessment of Summer and Autumn Carrots from a Biodynamic Breeding Project and Correlations of Physico-Chemical Parameters and Features Determined by Picture Forming Methods

*Ingo Hagel*<sup>1</sup>, *Dietrich Bauer*<sup>2</sup>, *Silvia Haneklaus*<sup>3</sup>, *Ewald Schnug*<sup>3</sup>

<sup>1</sup> *Institut für Biologisch-Dynamische Forschung, Brandschneise 5, D-64295 Darmstadt*

<sup>2</sup> *Dottenfelderhof, D-61118 Bad Vilbel*

<sup>3</sup> *Bundesforschungsanstalt für Landwirtschaft, Bundesallee 50, D-38116 Braunschweig*  
*eMail of the corresponding author: Ingo.Hagel@IBDF.de*

Keywords: *Breeding, Carrot, Picture Forming Methods, Quality*

### Introduction

Assessment of product quality is of special significance in organic farming and includes the supervision of crop quality in different growing systems (e.g., Fleck *et al.* 1998) and the characterisation of different cultivars. Several methods have been developed and applied for this purpose, e.g., the physico-chemical analysis of crops, picture forming methods (PFMs) and plant observation. So far only limited information is available on the comparability of these methods. This contribution aims to compare the results of the analysis of physico-chemical parameters of summer and autumn carrots with features determined by PFMs by means of correlation analysis.

### Material and Methods

Six summer and eight autumn carrot varieties from biodynamic breeders (D. Bauer, and Th. Heinze) and conventional breeders (Sperli, Quedlinburg) including open pollinating cultivars, F1-Hybrids and special selections were tested in a biodynamic breeding project established in 1998. These field experiments were carried out without replicates. The following parameters were determined in the plant samples collected at harvest time: dry matter, single carrot weight of autumn varieties, essential plant nutrients (N, NO<sub>3</sub>, P, K, Mg, S, Ca) and different carbohydrates. Also a decomposition test according to Reinhold (1943) was carried out in order to assess storage quality (Samaras, 1978). Because of a misunderstanding between project partners single carrot weight was only determined for autumn carrots. Quality assessments applying PFMs such as copper chloride crystallisation and chromatography were carried out and quantified from Dr. U. Balzer-Graf (Balzer-Graf, 1999 a and b) according to Balzer-Graf (2000).

### Results and Discussion

This contribution aims to compare the two methods rather than discussing the individual results obtained. This information is provided by Hagel *et al.* (2000 a and b) and Balzer-Graf (1999 a and b). The physico-chemical parameters of summer and autumn carrots (table 1 and 2) differed widely as did the results obtained by PFMs (table 3 and 4).

Correlation coefficients of parameters determined by physico-chemical analysis and PFMs are presented in tables 5 and 6. Due to low number of variants per regression (summer carrots: n = 6; autumn carrots: n = 8) sometimes even high correlation coefficients failed to turn out significant (e.g., P/Vitality for summer carrots: r = 0.81; fructosis/embryo-like: r = -0.79 (Table 5)). In order to illustrate the potential of relations for future projects using more variants these correlation coefficients were also mentioned in tables 5 and 6.

For summer carrots in general all nutritional elements indicated close relationships to features obtained by PFMs. E.g., for "Ripeness" determined by PFMs and the contents of K, Mg, S and Ca very high correlation coefficients from 0.93\*\* to 0.97\*\* were found (table 5). The closest significant correlation was found for Mg/Differentiation and Mg/Regeneration with r = 0.98 \*\*\* (table 5). Of course the strong relationship for Dry Matter/Ripeness (r = 0.87\*) appears reasonable. Also the positive relation for % N/Ripeness (r = 0.90\*) is consistent with findings from Wistinghausen (1979), who carried out time series analyses of carrots. But interestingly not for summer carrots (but for autumn carrots!) any significant correlations between parameters from PFMs and "Nitrate" were found (table 5 and 6).

Also the individual carbohydrates were more often correlated with features determined by PFMs in autumn than summer carrots (table 5 and 6). The reason could be related to an increased translocation rate of carbohydrates to the roots (as measured by content of dry matter) in autumn carrots due to the longer vegetation period (table 1 and 2). On the other hand a less pronounced relationship between mineral elements and PFMs in autumn carrots compared to summer carrots was found. With increasing single carrot weight the autumn carrots were judged more "Fruit-like" (r = 0.95\*\*\*) and less "Vegetative" (r = -0.94\*\*\*) by PFMs (table 6).

**Table 1.:** Mean content of dry matter (DM), loss of DM (%) in the decomposition test (DCT), N in % FM<sup>1</sup> (N), nitrate in ppm FM (NO<sub>3</sub>), nutrient elements (ppm in FM), carbohydrates in FM (%): glucosis (GLU), fructosis (FRU), monosaccharides in (MS), saccharosis in (SAC), total sugar (TS), monosaccharides:disaccharides (Mo:Di) of summer carrots in a biodynamic breeding project.

| Summer Carrots                    | DM   | DCT | N    | NO <sub>3</sub> | P   | K    | Mg  | S   | Ca  | GLU  | FRU  | MS   | SAC  | TS   | Mo:Di |
|-----------------------------------|------|-----|------|-----------------|-----|------|-----|-----|-----|------|------|------|------|------|-------|
| MICHEL (Bauer <sup>2</sup> )      | 12.1 | 56  | 0.16 | 115             | 359 | 2333 | 131 | 226 | 469 | 1.67 | 1.30 | 2.97 | 2.83 | 5.79 | 1.05  |
| MICHEL (Sperli)                   | 10.5 | 53  | 0.12 | 155             | 273 | 1976 | 125 | 189 | 419 | 2.15 | 1.70 | 3.85 | 2.02 | 5.87 | 1.91  |
| FRÜHBUND (Bauer)                  | 12.1 | 51  | 0.14 | 156             | 259 | 2271 | 125 | 228 | 422 | 2.06 | 1.66 | 3.72 | 2.94 | 6.67 | 1.27  |
| FRÜHBUND (Sperli)                 | 10.8 | 46  | 0.12 | 186             | 258 | 1941 | 117 | 179 | 385 | 2.15 | 1.76 | 3.92 | 2.28 | 6.19 | 1.72  |
| Mean of OP <sup>3</sup> Cultivars | 11.4 | 52  | 0.14 | 153             | 287 | 2130 | 124 | 206 | 424 | 2.01 | 1.61 | 3.62 | 2.52 | 6.13 | 1.49  |
| ANGLIA F1                         | 9.8  | 49  | 0.11 | 137             | 229 | 1632 | 98  | 155 | 359 | 2.35 | 1.73 | 4.08 | 2.02 | 6.10 | 2.03  |
| NANDA F1                          | 11.0 | 48  | 0.12 | 128             | 249 | 1695 | 104 | 158 | 393 | 2.24 | 1.62 | 3.86 | 2.85 | 6.72 | 1.35  |
| Mean of Hybrids                   | 10,4 | 49  | 0,12 | 133             | 239 | 1663 | 101 | 157 | 376 | 2,30 | 1,68 | 3,98 | 2,44 | 6,41 | 1,69  |

**Table 2.:** Mean content of dry matter (DM), single carrot weight (SCW), % loss of DM in decomposition test (DCT), % N in FM<sup>1</sup> (N), ppm nitrate in FM (NO<sub>3</sub>), nutrient elements (ppm in FM), carbohydrates in FM (%): glucosis (GLU), fructosis (FRU), monosaccharides (MS), saccharosis (SAC), total sugar (TS), monosaccharides:disaccharides (Mo:Di) of autumn carrots from a biodynamic breeding project.

| Autumn Carrots                                 | DM   | SCW | DCT | N    | NO <sub>3</sub> | Ca  | K    | P   | S   | Mg  | GLU  | FRU  | MS   | SAC  | TS   | Mo:Di |
|--|------|-----|-----|------|-----------------|-----|------|-----|-----|-----|------|------|------|------|------|-------|
| RODELIKA 97 (Bauer <sup>2</sup> )              | 14.6 | 143 | 41  | 0.15 | 80              | 441 | 2295 | 329 | 191 | 131 | 0.60 | 0.51 | 1.11 | 7.47 | 8.58 | 0.15  |
| RODELIKA 97 NUSSIG (Bauer)                     | 12.7 | 165 | 37  | 0.15 | 91              | 381 | 2492 | 284 | 158 | 119 | 1.30 | 1.08 | 2.39 | 4.62 | 7.00 | 0.52  |
| ROBILA (Heinze)                                | 13.1 | 122 | 51  | 0.18 | 280             | 464 | 2590 | 315 | 219 | 154 | 0.65 | 0.66 | 1.31 | 5.45 | 6.76 | 0.42  |
| LANGE ROTE STUMPFE (Quedlinburg <sup>2</sup> ) | 12.2 | 118 | 52  | 0.17 | 235             | 442 | 2578 | 357 | 212 | 129 | 0.92 | 0.79 | 1.71 | 4.24 | 5.95 | 0.40  |
| LANGE ROTE STUMPFE (Bauer)                     | 11.8 | 108 | 55  | 0.15 | 369             | 467 | 2589 | 297 | 194 | 118 | 1.14 | 0.86 | 2.00 | 4.05 | 6.06 | 0.49  |
| DUWICKA  | 11.1 | 53  | 64  | 0.14 | 223             | 310 | 3158 | 334 | 138 | 120 | 1.70 | 1.71 | 3.41 | 2.81 | 6.22 | 1.21  |
| Mean of OP <sup>3</sup> Cultivars              | 12,6 | 118 | 50  | 0,16 | 213             | 418 | 261  | 319 | 185 | 129 | 1,05 | 0,94 | 1,99 | 4,77 | 6,76 | 0,53  |
| TINO F1  | 10.1 | 74  | 45  | 0.12 | 211             | 430 | 1421 | 202 | 164 | 130 | 1.37 | 1.07 | 2.44 | 3.22 | 5.67 | 0.76  |
| NEVIS F1                                       | 10.1 | 87  | 38  | 0.11 | 349             | 385 | 1901 | 241 | 145 | 94  | 1.40 | 1.11 | 2.50 | 2.93 | 5.43 | 0.85  |
| Mean of Hybrids                                | 10,1 | 67  | 42  | 0,12 | 280             | 408 | 1661 | 222 | 155 | 112 | 1,39 | 1,09 | 2,47 | 3,08 | 5,55 | 0,81  |

<sup>1</sup> FM fresh matter; <sup>2</sup> breeder, <sup>3</sup> OP = open pollinating

**Table 3.:** Quality parameters according to picture forming methods of summer carrots from a biodynamic breeding project. Index ranging from 0 (very low) to 100 (very high). (from Balzer-Graf, 1999)

| Summer Carrots                | Carrot typical | Differen-<br>tiation | Vitality | Ripe-<br>ness | Regene-<br>ration | Embryo-<br>like | Vege-<br>tative | Dying | Leaf-<br>like |
|-------------------------------|----------------|----------------------|----------|---------------|-------------------|-----------------|-----------------|-------|---------------|
| MICHEL (Bauer <sup>1</sup> )  | 60             | 50                   | 70       | 80            | 60                | 40              | 0               | 10    | 0             |
| MICHEL (Sperli <sup>1</sup> ) | 60             | 45                   | 55       | 60            | 40                | 0               | 0               | 20    | 0             |
| FRÜHBUND (Bauer)              | 50             | 45                   | 60       | 70            | 40                | 30              | 0               | 0     | 20            |
| FRÜHBUND (Sperli)             | 50             | 40                   | 45       | 50            | 30                | 0               | 10              | 0     | 10            |
| ANGLIA F1                     | 30             | 20                   | 20       | 20            | 0                 | 0               | 40              | 50    | 0             |
| NANDA F1                      | 40             | 20                   | 40       | 40            | 0                 | 0               | 40              | 40    | 0             |

<sup>1</sup> breeder

**Table 4.:** Quality parameters according to Picture Forming Methods of autumn carrots from a biodynamic breeding project. Index ranging from 0 (very low) to 100 (very high). (from Balzer-Graf, 1999)

| Autumn Carrots                                | Carrot typical | Differen-<br>tiation | Vitality | Fruit-<br>like | Root-<br>like | Stabi-<br>lity | Vege-<br>tative | Ageing |
|---|----------------|----------------------|----------|----------------|---------------|----------------|-----------------|--------|
| RODELIKA 97 (Bauer <sup>1</sup> )             | 80             | 80                   | 70       | 90             | 80            | 60             | 0               | 0      |
| RODELIKA 97 NUSSIG (Bauer)                    | 80             | 70                   | 70       | 80             | 80            | 60             | 0               | 20     |
| ROBILA (Heinze <sup>1</sup> )                 | 70             | 60                   | 70       | 60             | 60            | 50             | 0               | 20     |
| LRSt <sup>2</sup> (Quedlinburg <sup>1</sup> ) | 50             | 50                   | 60       | 50             | 60            | 50             | 20              | 20     |
| LRSt <sup>2</sup> (Bauer)                     | 40             | 40                   | 50       | 40             | 40            | 40             | 30              | 30     |
| DUWICKA                                       | 40             | 80                   | 60       | 0              | 60            | 50             | 60              | 20     |
| TINO F1                                       | 20             | 10                   | 30       | 20             | 20            | 20             | 60              | 60     |
| NEVIS F1                                      | 30             | 20                   | 40       | 20             | 20            | 30             | 60              | 60     |

<sup>1</sup> breeder; <sup>2</sup> LRSt = LANGE ROTE STUMPFE

**Table 5.:** Correlation coefficients for the relationships between parameters of physico-chemical analysis and picture forming methods of summer carrots in a biodynamic breeding project. Number of variants per regression: n = 6. Statistical significances: p < 0.05 (\*); p < 0.01 (\*\*); p < 0.001 (\*\*\*)

|             | Carrot-<br>typical | Differen-<br>tiation | Vitality | Ripe-<br>ness | Regene-<br>ration | Embryo-<br>like | Vege-<br>tative | Dying   | Leaf-<br>like |
|-------------|--------------------|----------------------|----------|---------------|-------------------|-----------------|-----------------|---------|---------------|
| DM          |                    | 0.65                 | 0.86 *   | 0.87 *        | 0.70              | 0.88 *          | -0.61           | -0.69   |               |
| DCT         | 0.63               | 0.63                 | 0.71     | 0.71          | 0.73              | 0.75            | -0.69           |         |               |
| N           | 0.69               | 0.76                 | 0.88 *   | 0.90 *        | 0.86 *            | 0.96 **         |                 | -0.60   |               |
| NO3         |                    |                      |          |               |                   |                 |                 |         |               |
| P           | 0.74 *             | 0.70                 | 0.81     | 0.80          | 0.82 *            | 0.76            | -0.60           |         |               |
| K           | 0.79               | 0.92 *               | 0.93 **  | 0.96 **       | 0.94 **           | 0.88 *          | -0.89 *         | -0.82 * |               |
| Mg          | 0.95 **            | 0.98 ***             | 0.96 **  | 0.97 **       | 0.98 ***          | 0.68            | -0.97 **        | -0.82 * |               |
| S           | 0.72               | 0.88 *               | 0.90 *   | 0.93 **       | 0.90 *            | 0.90 *          | -0.86 *         | -0.77   |               |
| Ca          | 0.84 *             | 0.80                 | 0.95 **  | 0.94 **       | 0.88 *            | 0.82 *          | -0.74           |         |               |
| Glucosis    | -0.75              | -0.77                | -0.88 *  | -0.88 *       | -0.87 *           | -0.87 *         | 0.68            |         |               |
| Fructosis   |                    |                      | -0.64    | -0.62         |                   | -0.79           |                 |         |               |
| Saccharosis |                    |                      |          |               |                   | 0.70            |                 |         |               |
| Total Sugar |                    |                      |          |               |                   |                 |                 |         |               |
| Mo:Di       |                    |                      | -0.71    | -0.70         |                   | -0.81           |                 |         |               |

Only correlation coefficients > ±0.50 are listed

Different characteristics between the two sets of carrot variants were also noticed with regard to relations between the decomposition test (DCT) and PFM parameters. Only summer carrots showed reasonably high (though not significant) correlation coefficients from 0.63 to 0.75 (table 5). In contrast for autumn carrots all correlation coefficients were below 0.50 (therefore not shown in table 6).

The very often high and significant correlations between physico-chemical parameters and features determined by PFMs are striking and underline the technical accuracy of both methods. The results clearly confirm PFMs as a suitable tool for quality assessment which provide reliable results. This is also corroborated by the investigations of Balzer-Graf (2000) who showed that it is possible to distinguish crops from different growing systems by PFMs.

**Table 6.:** Correlation for the relationship between parameters of physico-chemical analysis and picture forming methods of autumn carrots in a biodynamic breeding project.

Number of variants per regression: n = 8. Statistical significances: p < 0.05 (\*); p < 0.01 (\*\*); p < 0.001 (\*\*\*)

|             | Carrot typical | Differentiation | Vitality | Fruit-like | Root-like | Stability | Vegetative | Ageing   |
|-------------|----------------|-----------------|----------|------------|-----------|-----------|------------|----------|
| SCW         | 0.87 **        |                 | 0.68     | 0.95 ***   | 0.67      | 0.66      | -0.94 ***  | -0.55    |
| DM          | 0.92 **        | 0.75 *          | 0.86 **  | 0.88 **    | 0.86 **   | 0.84 *    | -0.91 **   | -0.90 ** |
| DCT         |                |                 |          |            |           |           |            |          |
| N           |                | 0.55            | 0.53     |            | 0.63      | 0.65      |            | -0.71 *  |
| NO3         | -0.78 **       | -0.61           | -0.68    | -0.71 *    | -0.74 *   | -0.63     | 0.68       | 0.60     |
| P           | 0.54           | 0.78 *          | 0.78 *   |            | 0.73 *    | 0.80 *    |            | -0.86 ** |
| K           |                | 0.67            | 0.73 *   |            | 0.54      | 0.61      |            | -0.62    |
| Mg          |                |                 |          |            |           |           |            |          |
| S           |                |                 |          | 0.55       |           |           | -0.60      |          |
| Ca          |                |                 |          |            |           |           |            |          |
| Glucosis    | -0.59          |                 |          | -0.79 *    |           |           | 0.74*      | 0.55     |
| Fructosis   |                |                 |          | -0.68      |           |           | 0.58       |          |
| Saccharosis | 0.80 *         | 0.54            | 0.65     | 0.87 **    | 0.69      | 0.65      | -0.81 *    | -0.73 *  |
| Total Sugar | 0.86 **        | 0.74 *          | 0.75 *   | 0.79 *     | 0.80 *    | 0.76 *    | -0.76 *    | -0.81 *  |
| Mono:Di     | -0.65          |                 |          | -0.89 **   |           |           | 0.84 **    |          |

Only correlation coefficients > ±0.50 are listed

## Conclusions

High and significant correlation coefficients were found between quality parameters of summer and autumn carrots from a biodynamical breeding project determined by physico-chemical analysis and PFMs. This indicates close relationships between the two quality approaches which should be investigated further in future work.

## References

- Balzer-Graf, U. (1987): Vitalaktivität von Nahrungsmitteln im Spiegel bildschaffender Methoden. Elemente der Naturwissenschaft, Nr.46, 69-92.
- Balzer-Graf, U. (1999 a): Vitalqualität von Populations- und Hybridsorten bei Frühmöhren. Bericht zu Untersuchungen im Auftrag von Dietrich Bauer.
- Balzer-Graf, U. (1999 b): Vitalqualität von Populations- und Hybridsorten bei Lagermöhren. Bericht zu Untersuchungen im Auftrag von Dietrich Bauer.
- Balzer-Graf, U. (2000): Vitalquality. Quality Research with Picture Forming Methods. In: Lockeretz, W. and Geier, B. (Eds.), Quality and Communications for the organic Market, IFOAM, 179-188.
- Fleck, M., I. Hagel und A. Meier-Ploeger (1998): Lagerfähigkeit und Inhaltsstoffe von Möhren aus der biologisch-dynamischen und konventionellen Praxis. Deutsche Gesellschaft für Qualitätsforschung (Pflanzliche Nahrungsmittel) e.V., XXXIII. Vortragstagung, 23./24. März 1998, Dresden, 217-222.
- Hagel, I., D. Bauer, S. Haneklaus und E. Schnug (2000): Zur Qualität von Frühmöhren aus einem biologisch-dynamischen Züchtungsprojekt. Deutsche Gesellschaft für Qualitätsforschung e.V., XXXV. Vortragstagung, 20./21. März 2000, Karlsruhe (im Druck).
- Hagel, I., D. Bauer, S. Haneklaus und E. Schnug (2000): Zur Qualität von Herbstmöhren aus einem biologisch-dynamischen Züchtungsprojekt. Deutsche Gesellschaft für Qualitätsforschung e.V., XXXV. Vortragstagung, 20./21. März 2000, Karlsruhe (im Druck).
- Reinhold, J. (1943): Die Haltbarkeit von Gemüse, ein wichtiger Qualitätsfaktor. Z. Pflanzenkrankh. PflSchutz 53, 175-199.
- Samaras, I. (1978): Nachernteverhalten unterschiedlich gedüngter Gemüsearten mit besonderer Berücksichtigung physiologischer und mikrobiologischer Parameter. Diss., Gießen.
- WISTINGHAUSEN, E. v. (1979): Was ist Qualität? Wie entsteht sie und wie ist sie nachzuweisen? Verlag "Lebendige Erde", Darmstadt.

## **Bibliographische Angaben zu diesem Dokument:**

Hagel, I. und Bauer, D. und Haneklaus, S. und Schnug, E. (2000) Quality Assessment of Summer and Autumn Carrots from a Biodynamic Breeding Project and Correlations of Physico-Chemical Parameters and Features Determined by Picture Forming Methods. Beitrag präsentiert bei der Konferenz: 13th IFOAM Scientific Conference, Basel, 28.08.2000 - 31.08.2000; Veröffentlicht in Alföldi, Thomas und Lockeretz, William und Niggli, Urs, (Hrsg.) *Proceedings 13th International IFOAM Scientific Conference*, Seite(n) 284-287. IFOAM.

Das Dokument ist in der Datenbank „Organic Eprints“ archiviert und kann im Internet unter <http://orgprints.org/00002191> abgerufen werden.